

Airborne InSAR - 1974



From Graham, 1974

Optical Radar Correlator


Apollo Lunar Sounder Experiment




From Porcello et al., 1974

Airborne systems for measuring topography (1986)







SAR Processing Algorithms




- Many algorithms: Range-Doppler, Wave Domain, Omega-K, Spotlight, Spectral Analysis, Chirp Scaling, Stolt Interpolation
- But it's all *matched filtering*




COSEISMIC INTERFEROGRAMS AND GPS DISPLACEMENTS ICELAND EARTHQUAKES ON JUNE 17 AND 21, 2000







Magellan Mission to Venus



Burst Mode Pulse Encoding



ScanSAR Imaging Method and Geometry



SRTM Topography with Landsat Color Santa Barbara, CA




Bam Earthquake, Iran

Envisat Wide swath - Wide swath Interferogram

Baseline: 100 m


Burst overlap: 84%

(produced by polimi, courtesy Andrea Monti-Guarneri)




← ~400 km — →

Persistent Scatterers - Principle



Time and Varying Baseline

San Francisco Bay Area, CA 115,487 persistent scattering data points



From: Ferretti et al., EOS, 24 Aug 2004

Key:


Data acquired 1992-2000

SAF, HF, CF: San Andreas, Hayward, and Calaveras faults

CB Cupertino Basin, SL San Leandro Basin, AL Alameda, TI Treasure Island, BH Berkeley Hills


Persistent Scatterer Landslide Detail

Berkeley Hills, CA




From: Hilley et al., Science, 25 Jun 2004

Time / Rainfall Dependence of Landslide Area




Hilley et al., Science, 2004

Long Valley Caldera Observed with Persistent Scatterers



Time series InSAR over Long Valley caldera

Sept. 1992



Persistent scatterers

"Poor" quality interferogram "Good" quality interferogram Interferogram Persistent scatterers

Long Valley Caldera Unwrapped Time Series and Inferred Vertical Deformation

